

Note to Readers: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Desert Dust Outbreaks in Southern Europe: Contribution to Daily PM₁₀ Concentrations and Short-Term Associations with Mortality and Hospital Admissions

Massimo Stafoggia, Stefano Zauli-Sajani, Jorge Pey, Evangelia Samoli, Ester Alessandrini, Xavier Basagaña, Achille Cernigliaro, Monica Chiusolo, Moreno Demaria, Julio Díaz, Annunziata Faustini, Klea Katsouyanni, Apostolos G. Kelessis, Cristina Linares, Stefano Marchesi, Sylvia Medina, Paolo Pandolfi, Noemí Pérez, Xavier Querol, Giorgia Randi, Andrea Ranzi, Aurelio Tobias, Francesco Forastiere, and the MED-PARTICLES Study Group

Table of contents

Figure S1. Relative frequencies of desert dust days in the eleven cities of the MED-PARTICLES project, by season. Cities ordered by latitude, North to South. Desert dust days are defined as days with desert dust advection and estimated PM₁₀ concentrations at ground level > 0 µg/m³

Table S1. Daily PM concentrations (mean ± SD, µg/m³) by desert dust advection, in the eleven cities of the MED-PARTICLES project

Table S2. City-specific Pearson correlation coefficients between environmental variables. The lower triangle includes the correlations on days with desert dust. The upper triangle includes the correlations on days without desert dust. The cities are ordered by latitude, North to South. PM and ozone are measured as µg/m³, air temperature as °C

Table S2. City-specific Pearson correlation coefficients between environmental variables. The lower triangle includes the correlations on days with desert dust. The upper triangle includes the

correlations on days without desert dust. The cities are ordered by latitude, North to South. PM and ozone are measured as $\mu\text{g}/\text{m}^3$, air temperature as $^{\circ}\text{C}$ (continued)

Table S2. City-specific Pearson correlation coefficients between environmental variables. The lower triangle includes the correlations on days with desert dust. The upper triangle includes the correlations on days without desert dust. The cities are ordered by latitude, North to South. PM and ozone are measured as $\mu\text{g}/\text{m}^3$, air temperature as $^{\circ}\text{C}$ (continued)

Table S3. Daily mean counts of cause-specific mortality and hospital admissions in the eleven cities of the MED-PARTICLES project

Figure S2. Concentration–response relationship between non-desert PM_{10} (A) and desert PM_{10} (B) with natural mortality, lag 0-1. Values are estimated percent increases (95% CI) of total natural mortality associated with increases of PM_{10} levels relative to $5 \mu\text{g}/\text{m}^3$

Table S4. P-values of the city-specific likelihood-ratio tests comparing the models with natural spline and linear term for non-desert PM_{10} and desert PM_{10} exposures versus natural mortality